

Crepant Resolutions of $\mathbb{C}^n/A_1(n)$ and Flops of n -Folds for $n = 4, 5$

Li Chiang

Institute of Mathematics

Academia Sinica

Taipei , Taiwan

(e-mail:chiangl@gate.sinica.edu.tw)

Shi-shyr Roan

Institute of Mathematics

Academia Sinica

Taipei , Taiwan

(e-mail:maroan@ccvax.sinica.edu.tw)

Abstract

In this article, we determine the explicit toric variety structure of $\text{Hilb}^{A_1(n)}(\mathbb{C}^n)$ for $n = 4, 5$, where $A_1(n)$ is the special diagonal group of all order 2 elements. Through the toric data of $\text{Hilb}^{A_1(n)}(\mathbb{C}^n)$, we obtain certain toric crepant resolutions of $\mathbb{C}^n/A_1(n)$, and the different crepant resolutions are connected by flops of n -folds for $n = 4, 5$.

2000 MSC Primary: 14M25, 14J17, 20C33; Secondary: 13P10

1 Introduction

This article is intended to discuss the construction of flops of high-dimensional manifolds through the geometry of orbifolds. By an orbifold, we mean an algebraic (or analytic) variety with at most quotient singularities. This geometrical object was introduced in the 1950s by I. Satake, which he called a V-manifold, on his course of study of the Siegel's modular forms [22]. Special emphasis in this paper will be on the resolutions of orbifolds with the Gorenstein quotient singularities, i.e., the local structure around a singular point is given by

$$\mathbb{C}^n/G, \quad G : \text{a (nontrivial) subgroup of } \mathrm{SL}_n(\mathbb{C}), \quad n \geq 2. \quad (1)$$

In the mid-80s, string theorists studied a special type of physical model where strings propagate on an orbifold M/G where M is a manifold quotiented by a finite group G . The required string vacuum is described by the following orbifold Euler characteristic [5]:

$$\chi(M, G) = \frac{1}{|G|} \sum_{g, h \in G, gh=hg} \chi(M^{g,h})$$

where $M^{g,h}$ is the simultaneous fixed-point set of g, h . An equivalent expression of the above formula is given by

$$\chi(M, G) = \sum_{[g]} \chi(M^g/C(g)),$$

where $[g]$ is the conjugacy class of g with the centralizer $C(g)$. In the local situation (1), one obtains

$$\chi(\mathbb{C}^n, G) = |\mathrm{Irr}(G)|, \quad \mathrm{Irr}(G) := \{\text{irreducible representations of } G\}.$$

For the consistency of physical theory, the Gorenstein orbifold $\mathbb{C}^n/G$ demands a resolution $\widehat{\mathbb{C}^n/G}$ with the trivial canonical bundle and the Euler number $\chi(\widehat{\mathbb{C}^n/G}) = \chi(\mathbb{C}^n, G)$. For $n = 2$, all finite subgroups G of $\mathrm{SL}_2(\mathbb{C})$ were classified into *A-D-E* series by F. Klein, where the orbifolds $\mathbb{C}^2/G$ are always of the hypersurface singularity. The minimal resolution of $\mathbb{C}^2/G$ in the surface theory provides the required crepant resolution $\widehat{\mathbb{C}^2/G}$ [6]. For $n = 3$, the desired crepant resolution $\widehat{\mathbb{C}^3/G}$ was found for all G in the mid-nineties [7, 13, 20, 21] through some quantitative, but constructive, methods, heavily depending on the classical Miller-Blichfeldt-Dickson classification of finite subgroups of $\mathrm{SL}_3(\mathbb{C})$ [14], and the Klein's invariant theory of the simple groups I_{60} (Icosahedral group) and H_{168} (the simple group of order 168) [20]. The non-uniqueness of such crepant resolution is expected due to the flop-relation in 3-folds, but all with the required Euler number. Thereafter, the qualitative understanding and a possible canonical one of these crepant resolutions naturally arise as problems in algebraic geometry. For this purpose, the development has resulted in the concept of Hilbert scheme of G -orbits (or G -Hilbert scheme), $\mathrm{Hilb}^G(\mathbb{C}^n)$, associated to $\mathbb{C}^n/G$, which was proposed in [9] by engaging the finite-group representation theory into geometrical problems of orbifolds. The structure of G -Hilbert scheme has now provided a positive answer for the problem addressed to the crepant resolution with the orbifold Euler number for the dimension $n = 2, 3$ [2, 8, 16].

For $n \geq 4$, very few results have been known on the crepant resolution problem of $\mathbb{C}^n/G$, and the structure of $\mathrm{Hilb}^G(\mathbb{C}^n)$, partly due to the lack of common conclusion for all G , as $\mathbb{C}^n/G$ has no crepant resolution at all for a large class of groups G , even in the abelian group case. A suitable class of groups G , which could be of interest to study, are those with Gorenstein orbifold $\mathbb{C}^n/G$ of the *hypersurface type*, and one can regard it as one way to the high-dimensional generalization of Klein surface singularities. For the abelian group case, the group G is given by

$$A_r(n) := \{g \in \mathrm{SL}_n(\mathbb{C}) | g^{r+1} = 1, \text{ } g \text{ is diagonal}\} \quad , \quad r \geq 1.$$

In [3], we have studied the cases $A_r(4)$ for all r , and obtained the crepant resolutions $\widehat{\mathbb{C}^4/A_r(4)}$ through the detailed structure of $\text{Hilb}^{A_r(4)}(\mathbb{C}^4)$. While in this work, we shall discuss the cases $A_1(n)$ for $n = 4, 5$, in order to explore the flop-relation of n -folds. It is known that there do exist some crepant resolutions of $\mathbb{C}^n/A_1(n)$ for all n in [20], where the result was obtained by blowing-up techniques through the hypersurface equation of $\mathbb{C}^n/A_1(n)$. But in this paper, we shall construct another type of crepant resolutions of $\mathbb{C}^n/A_1(n)$, which are toric varieties, through the toric structure of $\text{Hilb}^{A_1(n)}(\mathbb{C}^n)$. One of the main goals of this approach is to identify the proper “flop” concept of n -folds for $n = 4, 5$. This method is consistent with the flops of 3-folds, considered as crepant resolutions of the 3-dimensional isolated singularity with the equation, $XY = ZW$, a construction known since the seventies in the study of degeneration of K3 surfaces [17]. However, for $n = 4, 5$, the singularities, which possess crepant resolutions with the flop relation discovered in this article, has a complicated expression of algebraic equations (see formula (5), (6) of the paper). In particular, it is not of the type of complete intersection.

This paper is organized as follows: In §2, we give a brief survey on the construction of crepant resolution $\widehat{\mathbb{C}^n/G}$ for $n = 2, 3$. In §3, we review some basic facts in G -Hilbert schemes and toric geometry for later discussions. In §4, we derive the toric variety structure of $\text{Hilb}^{A_1(4)}(\mathbb{C}^4)$, which is smooth but not crepant. By blowing down the canonical divisor $(\mathbb{P}^1)^3$ of $\text{Hilb}^{A_1(4)}(\mathbb{C}^4)$ to $(\mathbb{P}^1)^2$ in different factors, we obtain three toric crepant resolutions of $\mathbb{C}^4/A_1(4)$. By this process, the flop of 4-folds is naturally revealed between these three crepant resolutions. In §5, we derive the toric variety structure of $\text{Hilb}^{A_1(5)}(\mathbb{C}^5)$, which is singular and non-crepant. By analyzing the toric structure of $\text{Hilb}^{A_1(5)}(\mathbb{C}^5)$, we construct twelve crepant resolutions of $\mathbb{C}^5/A_1(5)$, all dominated by $\text{Hilb}^{A_1(5)}(\mathbb{C}^5)$. The connection between these crepant resolutions gives rise to the “flop” relation of 5-folds.

2 Kleinian Singularities and Crepant Resolutions of Gorenstein Three-Dimensional Orbifolds

For a finite non-trivial subgroup G of $\text{SL}_2(\mathbb{C})$, $\mathbb{C}^2/G$ is a surface with an isolated singularity at the origin. These singularities were classified by Klein in 1872 (or so) in his work on the invariant theory of regular solids in $\mathbb{R}^3$, and they are described in the following table:

Type	Group G	$ G $	degree of invariants	relation of invariants
A_r	Abelian	$r + 1$	$2, r + 1, r + 1$	$X^{r+1} + YZ$
D_r	Binary dihedral	$4(r - 2)$	$4, 2(r - 2), 2(r - 1)$	$X^{r-1} + XY^2 + Z^2$
E_6	Binary tetrahedral	24	6, 8, 12	$X^3 + Y^4 + Z^2$
E_7	Binary octahedral	48	8, 12, 18	$X^3Y + Y^3 + Z^2$
E_8	Binary icosahedral	120	12, 20, 30	$X^5 + Y^3 + Z^2$

For A -type group G , it is a cyclic subgroup of $\text{SL}_2(\mathbb{C})$ generated by the diagonal element with the entries $e^{\frac{2\pi i}{r+1}}, e^{\frac{2\pi ir}{r+1}}$. By the continued fraction,

$$\frac{r+1}{r} = 2 - \frac{1}{2 - \frac{1}{\ddots}},$$

the minimal resolution $\widehat{\mathbb{C}^2/G}$ has the exceptional set consisting of a tree of r -rational curves with self-intersections -2 , which implies the trivial canonical bundle of $\widehat{\mathbb{C}^2/G}$ with the Euler number $r + 1$. For other groups G in $\text{SL}_2(\mathbb{C})$, the minus identity map always belongs to the center of G . One can construct the minimal resolution by first blowing up $\mathbb{C}^2$ at the origin, then analyzing the fixed-points of G on this blowing-up space, and then reducing to cyclic quotient singularities. The following remarkable characterization for Kleinian singularities by Brieskorn is known:

THEOREM 2.1 [1] *For an isolated normal surface singularity, the following are equivalent: (1) it is Kleinian, (2) it is rational with multiplicity 2, (3) it is rational with embedding dimension 3, and (4) the minimal resolution has the exceptional set by a configuration of rational (-2)-curves with an intersection matrix minus that of the Cartan matrix of the root system of type A_r, D_r, E_6, E_7 or E_8 . The correspondence between (1) and (4) is indicated in the above table.*

On the other hand, J. McKay observed an interesting connection between the representation theory of Kleinian groups and the affine Dynkin diagrams of the A - D - E root system. Denote $c : G \longrightarrow \mathrm{GL}_2(\mathbb{C})$ the ‘canonical’ representation of G . For $\rho, \rho' \in \mathrm{Irr}(G)$, let $m(\rho, \rho')$ be the multiplicity of ρ' in the representation $\rho \otimes c$:

$$\rho \otimes c = \sum_{\rho' \in \mathrm{Irr}(G)} m(\rho, \rho') \rho'.$$

By the self-dual property of c , $m(\rho, \rho') = m(\rho', \rho)$ for all ρ, ρ' . One forms a diagram $\tilde{\Gamma}(G)$ with elements in $\mathrm{Irr}(G)$ as vertices, and two vertices ρ, ρ' connected by $m(\rho, \rho')$ edges. Then, one has the following result:

THEOREM 2.2 [15] *$\tilde{\Gamma}(G)$ is isomorphic to the affine Dynkin diagram of the corresponding root system.*

By a suitably chosen isomorphism between $\tilde{\Gamma}(G)$ and the affine Dynkin diagram, the trivial representation of G corresponds to the negative longest root of the affine diagram. So, the diagram $\Gamma(G)$ obtained by deleting the trivial representation is isomorphic to the ordinary Dynkin diagram of its corresponding system. With the results in Theorems 2.1 and 2.2, the rational (-2)-curves in resolution configuration implies the trivial canonical bundle of $\widehat{\mathbb{C}^2/G}$, and the Euler number equality follows from McKay’s correspondence [6].

For $n = 3$, all finite subgroups G of $\mathrm{SL}_3(\mathbb{C})$ were classified by Miller et al. in 1916 [14]. The list consists of 12 types of such groups, among which two well-known simple groups, I_{60} and H_{168} appear. For the problem of the crepant resolutions of $\widehat{\mathbb{C}^3/G}$, due to the long list of the Miller-Blichfeldt-Dickson classification, it is still hard to study the problem by the method of case by case approach. Hence, certain induction procedures are introduced, as the one of blowing up at the origin in the 2-dimensional case, to enable us to work only on a few cases, consisting of abelian groups, certain solvable groups and simple groups I_{60}, H_{168} [20]. For abelian groups, the toric geometry [10] provides the effective computational methods to obtain the solutions [12, 18, 19]. Using these techniques, one can work on an explicitly given solvable subgroup of $\mathrm{SL}_3(\mathbb{C})$ and obtain the crepant resolutions by using a certain ‘canonical’ procedure, regardless the resulting structure of the resolution might be complicated. For the simple groups I_{60}, H_{168} in $\mathrm{SL}_3(\mathbb{C})$, one needs some different techniques, using the invariant theory of their 3-dimensional representations in Klein’s work [11]:

G	degree of invariants	relation of invariants
I_{60}	2, 6, 10, 15	$-144W^2 + Z^3 - 1728Y^5 + 720XY^3Z$ $-80X^2YZ^2 + 64X^3(5Y^2 - XZ)^2$
H_{168}	4, 6, 14, 21	$-W^2 + Z^3 + 1728Y^7 + 1008XY^4Z$ $-88X^2YZ^2 - 60032X^3Y^5 + 1088X^4Y^2Z$ $+22016X^6Y^3 - 256X^7Z - 2048X^9Y$

Note that the relation of invariants in the above table define a hypersurface of $\mathbb{C}^3/G$ in $\mathbb{C}^4$. One can obtain the required crepant resolution by making full use of the relations of invariants of these simple

groups G [13, 20]. With all these efforts, one arrives at a solution of the crepant resolutions $\widehat{\mathbb{C}^3/G}$ for a finite subgroup G of $\mathrm{SL}_3(\mathbb{C})$ [21]. However, all of those works were carried out using an explicit (somewhat unpleasant) computational method, which is cumbersome and not very illuminating. An approach of a qualitative nature to the crepant resolution $\widehat{\mathbb{C}^3/G}$ is provided by the method of G -Hilbert scheme, which will be briefly described in the next section.

3 G -Hilbert scheme, Abelian Orbifolds and Toric Geometry

From now on for the rest of this paper, $\{e^1, \dots, e^n\}$ will always denote the standard basis of $\mathbb{C}^n$, $\{Z_1, \dots, Z_n\}$ the corresponding dual basis, and $\mathbb{C}[Z] := \mathbb{C}[Z_1, \dots, Z_n]$ the polynomial ring of Z_i 's.

For a finite subgroup G of $\mathrm{GL}_n(\mathbb{C})$, we denote $S_G = \mathbb{C}^n/G$ with the canonical projection $\pi_G : \mathbb{C}^n \rightarrow S_G$, and $o = \pi_G(\vec{0})$. In this article, a variety X is said to be birational over S_G if there exists a proper birational epimorphism $\sigma_X : X \rightarrow S_G$. In this case, we have the commutative diagram:

$$\begin{array}{ccc} X \times_{S_G} \mathbb{C}^n & \longrightarrow & \mathbb{C}^n \\ \downarrow \pi & & \downarrow \pi_G \\ X & \xrightarrow{\sigma_X} & S_G . \end{array}$$

Denote $\mathcal{F}_X (= \pi_* \mathcal{O}_{X \times_{S_G} \mathbb{C}^n})$ the coherent $\mathcal{O}_X$ -sheaf over X obtained by the push-forward of the structure sheaf of $X \times_{S_G} \mathbb{C}^n$. The geometrical fiber of $\mathcal{F}_X$ over an element y of X is given by $\mathcal{F}_{X,y} = k(y) \otimes_{\mathcal{O}_X} \mathcal{F}_X$, which is isomorphic to $\mathbb{C}[Z]/I(y)$ for some G -invariant ideal $I(y)$ in $\mathbb{C}[Z]$. The G -Hilbert scheme $\mathrm{Hilb}^G(\mathbb{C}^n)$ with the birational morphism, $\sigma_{\mathrm{Hilb}} : \mathrm{Hilb}^G(\mathbb{C}^n) \rightarrow S_G$, is the minimal object in the category of varieties X birational over S_G such that $\mathcal{F}_X$ is a vector bundle over X [9] (or see [3] and references therein). Furthermore, for each X in the category, there exists a unique epimorphism λ_X from X to $\mathrm{Hilb}^G(\mathbb{C}^n)$. Indeed, $\lambda_X(y)$ is represented by the ideal $I(y)$. In the case when $\mathbb{C}^n/G$ in (1), for $n = 2$, $\mathrm{Hilb}^G(\mathbb{C}^2)$ is the minimal resolution of $\mathbb{C}^2/G$, and for $n = 3$, $\mathrm{Hilb}^G(\mathbb{C}^3)$ is a toric crepant resolution of $\mathbb{C}^3/G$ for an abelian group G [8, 16]. For a general finite subgroup G of $\mathrm{SL}_3(\mathbb{C})$, $\mathrm{Hilb}^G(\mathbb{C}^3)$ is indeed a crepant resolution $\widehat{\mathbb{C}^3/G}$ has been justified in [2] by methods in homological algebra, an argument bypassing the geometry of G -Hilbert scheme (of which explicit structures are important on its own from the aspect of applications to certain relevant physical problems).

When G is an abelian group, one can apply toric geometry to study problems on resolutions of S_G . We now give a brief review of some facts in toric geometry for later use in this article (for the details, see [10]). For the rest of this paper, we shall always assume G to be a finite abelian group in $\mathrm{SL}_n(\mathbb{C})$, and identify G as a subgroup of the diagonal group $T_0 := \mathbb{C}^{*n} \subset \mathrm{GL}_n(\mathbb{C})$. Regarding $\mathbb{C}^n$ as the partial compactification of T_0 , we define the n -torus T with the embedding in the T -space S_G by

$$T := T_0/G, \quad T \subset S_G .$$

Denote $N = \mathrm{Hom}(\mathbb{C}^*, T)$ (resp. $N_0 = \mathrm{Hom}(\mathbb{C}^*, T_0)$) the lattice of one-parameter subgroups of T (resp. T_0), M (resp. M_0) the dual lattice of N (resp. N_0). We have $N \supset N_0$ and $M \subset M_0$. Through the map $\exp : \mathbb{R}^n \rightarrow T_0$, $\exp(\sum_i x_i e^i) := \sum_i e^{2\pi\sqrt{-1}x_i} e^i$, one can identify N_0 and N with the following lattices in $\mathbb{R}^n$,

$$N_0 = \mathbb{Z}^n (= \exp^{-1}(1)) , \quad N = \exp^{-1}(G) .$$

In this way, M_0 can be identified with the monomial group of variables $Z_1, \dots, Z_n$, and M corresponds to the subgroup of G -invariant monomials in M_0 .

By [10], a T -space birational over S_G is described by a fan $\Sigma = \{\sigma_l \mid l \in I\}$ with $C_0 := \sum_i \mathbb{R}_{\geq 0} e^i$ as its support, i.e., a rational polyhedral cone decomposition of C_0 . In this situation, an equivalent

description is given by the polytope decomposition $\Lambda = \{\Delta_l \mid l \in I\}$ of the simplex

$$\Delta := \left\{ \sum_i x_i e^i \in C_0 \mid \sum_i x_i = 1 \right\},$$

where $\Delta_l := \sigma_l \cap \Delta$ with the vertices of Δ_l in $\Delta \cap \mathbb{Q}^n$. For $\sigma_l = \{\mathbf{0}\}$, we have $\Delta_l = \emptyset$. We shall call Λ a rational polytope decomposition of Δ , and denote X_Λ the toric variety corresponding to Λ .

For a rational polytope decomposition Λ of Δ , we define $\Lambda(i) := \{\Delta_l \in \Lambda \mid \dim(\Delta_l) = i\}$ for $-1 \leq i \leq n-1$, (here $\dim(\emptyset) := -1$). Then, for each $\Delta_l \in \Lambda(i)$, there associates a T -orbit of dimension $n-1-i$, denoted by $\text{orb}(\Delta_l)$. For $\Delta_l \in \Lambda(n-1)$, i.e., Δ_l is an $(n-1)$ -dimensional polytope, $\text{orb}(\Delta_l)$ consists of only one point, which will be denoted by x_{Δ_l} . For a vertex $v \in \Lambda(0)$, the closure of $\text{orb}(v)$ represents a toric divisor, which will be denoted by D_v . The canonical sheaf of X_Λ is given by the following expression of toric divisors,

$$\omega_{X_\Lambda} = \mathcal{O}_{X_\Lambda} \left(\sum_{v \in \Lambda(0)} (m_v - 1) D_v \right), \quad (2)$$

where m_v is the smallest positive integer with $m_v v \in N$, i.e., a primitive element of N .

A polytope decomposition Λ of Δ is called integral if all vertices of Λ are in N . By (2), the trivial canonical sheaf is described by the integral condition of Λ . The non-singular criterion of X_Λ is given by the simplicial decomposition of Λ with the multiplicity-one property. For a polytope $\mathcal{R} \in \Lambda(i)$, we denote

$$X_{\mathcal{R}} = \text{spec}(\mathbb{C}[M \cap \sigma_{\mathcal{R}}])$$

where $\sigma_{\mathcal{R}}$ is the dual of the cone $\sigma_{\mathcal{R}} := \{rv \mid r \geq 0, v \in \mathcal{R}\}$. Then, $\{X_{\mathcal{R}}\}_{\mathcal{R} \in \Lambda(n-1)}$ forms an open cover of X_Λ .

Now we consider the case $G = A_1(n)$. We shall always denote the following elements in $M (= M_0^G)$:

$$T_i = Z_i^2, \quad U_i = \frac{Z_i}{(Z_1 \dots \hat{Z}_j \dots Z_n)}, \quad V_{ij} = \frac{Z_i Z_j}{\prod_{\alpha \neq i, j; 1 \leq \alpha \leq n} Z_\alpha}. \quad (3)$$

The set of N -integral elements in Δ is given by

$$\Delta \cap N = \{e^i \mid 1 \leq i \leq n\} \cup \{v^{ij} \mid 1 \leq i < j \leq n\}, \quad v^{ij} := \frac{1}{2}(e^i + e^j).$$

Other than the simplex Δ itself, there is only one integral polytope decomposition of Δ , denoted by Ξ , which is invariant under all permutations of the coordinates (i.e., $\tau : e^i \mapsto e^{\tau(i)}$ for $\tau \in S_n$, the symmetric group of degree n). Then, $\Xi(n-1)$ consists of the following $n+1$ polytopes:

$$\diamond := \langle v^{ik} \mid 1 \leq i < k \leq n \rangle, \quad \Delta_j := \langle e^j, v^{ij} \mid i \neq j \rangle \quad (j = 1, \dots, n). \quad (4)$$

The toric variety X_Ξ is smooth on the affine space centered at x_{Δ_j} with the local coordinate U_j, T_i ($i \neq j$). Therefore, the singular set of X_Ξ lies in the affine open set $X_\diamond$ centered at $x_\diamond$. The structure near $x_\diamond$ is determined by $\check{\sigma}_\diamond \cap M$, which is generated by T_i, U_i^{-1} . For $n = 3$, $U_1^{-1}, U_2^{-1}, U_3^{-1}$ form an integral basis for M , which implies the smoothness of X_Ξ . For $n \geq 4$, X_Ξ is singular at $x_\diamond$, and $T_i, U_i^{-1}, 1 \leq i \leq n$, form the minimal generating set of $\check{\sigma}_\diamond \cap M$. For $n = 4$, the singular structure of X_Ξ near $x_\diamond$ is described by the 4-dimensional affine variety in $\mathbb{C}^8$ with the equations:

$$t_i \bar{u}_i = t_j \bar{u}_j, \quad t_i t_j = \bar{u}_i \bar{u}_j, \quad (t_i, \bar{u}_i)_{1 \leq i \leq 4} \in \mathbb{C}^8 \quad (5)$$

where $i \neq j$ and $\{i', j'\}$ is the complementary pair of $\{i, j\}$. For $n = 5$, the singular structure of X_Ξ near $x_\diamond$ is defined by the following relations:

$$t_i \bar{u}_i = t_j \bar{u}_j, \quad \bar{u}_i \bar{u}_j = t_k t_l t_m, \quad (t_i, \bar{u}_i)_{1 \leq i \leq 5} \in \mathbb{C}^{10} \quad (6)$$

where $1 \leq i < j \leq 5$, and $\{k, l, m\}$ is the complement of $\{i, j\}$.

For later use, we recall some terminology of the Gröbner basis [4]. Let $\prec_L$ be a lexicographic order on $\mathbb{C}[Z]$ and $\mathbf{w} \in \mathbb{Z}_{\geq 0}^n$. The weight order $\prec$ determined by the weight $\mathbf{w}$ is the monomial ordering on $\mathbb{C}[Z]$ with the following properties:

- (1) If $(\mathbf{u}_1 - \mathbf{u}_2) \cdot \mathbf{w} < 0$, then $Z^{\mathbf{u}_1} \prec Z^{\mathbf{u}_2}$.
- (2) If $(\mathbf{u}_1 - \mathbf{u}_2) \cdot \mathbf{w} = 0$ and $Z^{\mathbf{u}_1} \prec_L Z^{\mathbf{u}_2}$, then $Z^{\mathbf{u}_1} \prec Z^{\mathbf{u}_2}$.


For a monomial ideal $I \subset \mathbb{C}[Z]$, we shall denote $I^\perp$ the set of monic monomials outside I . For a monomial ordering $\prec$ and an ideal J , $\text{lt}_\prec(J)$ will denote the ideal generated by the leading terms (with respect to $\prec$) of all elements in J .

4 $\text{Hilb}^{A_1(4)}(\mathbb{C}^4)$ and Crepant Resolutions of $\mathbb{C}^4/A_1(4)$


In this section, we consider the case $n = 4$. Throughout this section, the indices j, k, l, m will always denote a permutation of $(1, 2, 3, 4)$. For $n = 4$, the polytope $\diamond$ in (4) is a regular 3-dimensional octahedron contained in the standard simplex Δ in $\mathbb{R}^4$, with a cube as its dual polygon. We label the facets of the octahedron $\diamond$ by F_j, F'_j where

$$F_j = \diamond \cap \Delta_j, \quad F'_j = \left\{ \sum_{i=1}^4 x_i e^i \in \diamond \mid x_j = 0 \right\}.$$

Then, the dual of F_j, F'_j are vertices of the cube, denoted by α_j, α'_j (see [Fig. 1]).



[Fig. 1] Dual pair of octahedron and cube: faces F_j, F'_j of octahedron are dual to vertices α_j, α'_j of cube.



[Fig. 2] The rational simplicial decomposition Ξ^* of Δ for $n = 4$.

Consider the rational simplicial decomposition Ξ^* of Δ obtained by Ξ and adding the center $c := \frac{1}{4} \sum_{j=1}^4 e^j$ as a vertex of Ξ^* with the barycentric decomposition of $\diamond$ in Ξ , (see [Fig. 2]). Then, $\Xi^*(3)$ consists of the polytopes Δ_j, C_j and C'_j for $j = 1, \dots, 4$, where

$$C_j := \langle c, v^{jk}, v^{jl}, v^{jm} \rangle, \quad C'_j := \langle c, v^{kl}, v^{lm}, v^{mk} \rangle.$$

Note that $c \notin N$ and $2c \in N$.

THEOREM 4.1 For $G = A_1(4)$, we have $\text{Hilb}^G(\mathbb{C}^n) \simeq X_{\Xi^*}$, which is non-singular with the canonical bundle $\omega = \mathcal{O}_{X_{\Xi^*}}(E)$, where E is an irreducible divisor isomorphic to the triple product of $\mathbb{P}^1$, $E = (\mathbb{P}^1)^3$. Furthermore, for $\{j, k, l\} = \{1, 2, 3\}$ and the (k, l) -th factor projection, $p_j : E \rightarrow (\mathbb{P}^1)^2$, the restriction of normal bundle of E on each fiber ($\simeq \mathbb{P}^1$) of p_j is the (-1) -hyperplane bundle of $\mathbb{P}^1$.

Proof: The smoothness of the affine spaces $X_{\mathcal{R}}$, $\mathcal{R} = \Delta_j, C_j, C'_j$, follows from the integral and multiplicity-one properties of $\sigma_{\mathcal{R}}$ with respect to N . With (3) for $n = 4$, we have $V_{jk} = (Z_j Z_k)/(Z_l Z_m)$, $U_j = Z_j/(Z_k Z_l Z_m)$. By using the inverse matrices of $(e^j, v^{jk}, v^{jl}, v^{jm})$, $(2c, v^{jk}, v^{jl}, v^{jm})$ and $(2c, v^{kl}, v^{lm}, v^{mk})$, one obtains the generators of $\sigma_{\mathcal{R}} \cap M$ for $\mathcal{R} = \Delta_j, C_j$, and C'_j as follows:

$$\begin{aligned} \Delta_j : & \quad U_j, \quad T_k, \quad T_l, \quad T_m; \\ C_j : & \quad U_j^{-1}, \quad V_{jk}, \quad V_{jl}, \quad V_{jm}; \\ C'_j : & \quad T_j, \quad V_{lm}, \quad V_{km}, \quad V_{kl}. \end{aligned}$$

We shall express the coordinates of an element $y \in X_{\mathcal{R}} \simeq \mathbb{C}^4$ as follows:

$$\begin{aligned} (U_j, T_k, T_l, T_m) &= (u_j, t_k, t_l, t_m), & \mathcal{R} &= \Delta_j, \\ (U_j^{-1}, V_{jk}, V_{jl}, V_{jm}) &= (\bar{u}_j, v_{jk}, v_{jl}, v_{jm}), & \mathcal{R} &= C_j, \\ (T_j, V_{lm}, V_{km}, V_{kl}) &= (t_j, v_{lm}, v_{km}, v_{kl}), & \mathcal{R} &= C'_j, \end{aligned} \tag{7}$$

and the corresponding ideal $I(y)$ is given by

$$\left. \begin{aligned} & \langle Z_j - u_j Z_k Z_l Z_m, Z_k^2 - t_k, Z_l^2 - t_l, Z_m^2 - t_m \rangle, & \mathcal{R} &= \Delta_j; \\ & \langle Z_k Z_l Z_m - \bar{u}_j Z_j, Z_j Z_k - v_{jk} Z_l Z_m, Z_j Z_l - v_{jl} Z_k Z_m, \\ & \quad Z_j Z_m - v_{jm} Z_k Z_l, Z_j^2 - t_j, Z_k^2 - t_k, Z_l^2 - t_l, Z_m^2 - t_m \rangle, & \mathcal{R} &= C_j; \\ & (t_j = \bar{u}_j v_{jk} v_{jl} v_{jm}, t_\alpha = \bar{u}_j v_{j\alpha}, (\alpha \neq j)), \\ & \langle Z_l Z_m - v_{lm} Z_j Z_k, Z_k Z_m - v_{km} Z_j Z_l, \\ & \quad Z_k Z_l - v_{kl} Z_j Z_m, Z_j^2 - t_j, Z_k^2 - t_k, Z_l^2 - t_l, Z_m^2 - t_m \rangle, & \mathcal{R} &= C'_j. \\ & (t_k = t_j v_{km} v_{kl}, t_l = t_j v_{lm} v_{kl}, t_m = t_j v_{km} v_{lm}), \end{aligned} \right\} \tag{8}$$

In each case, one can show that the generators in the expression of (8) form the reduced Gröbner basis of $I(y)$, and $\text{lt}_{\prec}(I(y)) = I(x_{\mathcal{R}})$ for a weight order $\prec$ with the weight in $\text{Interior}(\sigma_{\mathcal{R}})$. For $\mathcal{R} = \Delta_j$, we have $I(x_{\Delta_j}) = \langle Z_j, Z_k^2, Z_l^2, Z_m^2 \rangle$, and

$$I(x_{\Delta_j})^\perp = \{1, Z_k, Z_l, Z_m, Z_l Z_m, Z_k Z_m, Z_k Z_l, Z_k Z_l Z_m\}$$

gives rise to a G -regular monomial basis for $\mathbb{C}[Z]/I(y)$. Similarly, for $\mathcal{R} = C_j, C'_j$, we have $I(x_{C_j}) = \langle Z_k Z_l Z_m, Z_j Z_k, Z_j Z_l, Z_j Z_m, Z_1^2, Z_2^2, Z_3^2, Z_4^2 \rangle$ and $I(x_{C'_j}) = \langle Z_l Z_m, Z_k Z_m, Z_k Z_l, Z_1^2, Z_2^2, Z_3^2, Z_4^2 \rangle$ with

$$I(x_{C_j})^\perp = \{1, Z_1, Z_2, Z_3, Z_4, Z_l Z_m, Z_k Z_m, Z_k Z_l\},$$

$$I(x_{C'_j})^\perp = \{1, Z_1, Z_2, Z_3, Z_4, Z_j Z_k, Z_j Z_l, Z_j Z_m\},$$

which give rise to corresponding G -regular monomial basis of $\mathbb{C}[Z]/I(y)$. Since X_{Ξ^*} is birational over S_G , with the vector bundle $\mathcal{F}_{X_{\Xi^*}}$, we have an epimorphism

$$\lambda : X_{\Xi^*} \rightarrow \text{Hilb}^{A_1(4)}(\mathbb{C}^4)$$

with $I(\lambda(y)) = I(y)$ for $y \in X_{\Xi^*}$. We are going to show the injectivity of λ . For each $\mathcal{R} \in \Xi^*(3)$, the coordinates of an element in $X_{\mathcal{R}}$ can be represented in the form $(p_i/q_i = \gamma_i)_{i=1}^4$ for some monomials p_i, q_i described in (7). Let y, y' be two points in $X_{\mathcal{R}}$ with the coordinates $(p_i/q_i = \gamma_i)_{i=1}^4$ and $(p_i/q_i = \gamma'_i)_{i=1}^4$

respectively. By (7), (8), we have $p_i - \gamma_i q_i \in I(y)$ (resp. $p_i - \gamma'_i q_i \in I(y')$) with $q_i \in I(x_{\mathcal{R}})^\perp$ for each i . When $I(y') = I(y)$, we have $(\gamma'_i - \gamma_i)q_i \in I(y)$. By $q_i \in I(x_{\mathcal{R}})^\perp$, one has $\gamma'_i = \gamma_i$ for all i , which implies $y = y'$. Hence, λ is injective on each $X_{\mathcal{R}}$. Furthermore, the ideal $I(y)$ for $y \in X_{\mathcal{R}}$ is completely determined by its toric coordinates, which appear as some generating elements of $I(y)$ in (8). By the construction of toric variety, one can conclude that if $y \in X_{\mathcal{R}}$ and $y' \in X_{\mathcal{R}'}$ with $I(y) = I(y')$, the toric coordinates of y for $X_{\mathcal{R}}$ and y' for $X_{\mathcal{R}'}$ are related by the transition function of these affine charts, hence $y = y'$. For example, for $y \in X_{C'_4}$ and $y' \in X_{C_1}$, the affine coordinate of $y \in X_{C'_4}$ is given by

$$(\xi_1, \xi_2, \xi_3, \xi_4) := \left(Z_4^2, \frac{Z_2 Z_3}{Z_1 Z_4}, \frac{Z_1 Z_3}{Z_2 Z_4}, \frac{Z_1 Z_2}{Z_3 Z_4} \right)$$

and $y' \in X_{C_1}$ is

$$(\eta_1, \eta_2, \eta_3, \eta_4) := \left(\frac{Z_2 Z_3 Z_4}{Z_1}, \frac{Z_1 Z_4}{Z_2 Z_3}, \frac{Z_1 Z_3}{Z_2 Z_4}, \frac{Z_1 Z_2}{Z_3 Z_4} \right).$$

By the expression of ξ 's and η 's in terms of Z_i 's, we obtain the transition function on $X_{C'_4} \cap X_{C_1}$:

$$(\xi_1, \xi_2, \xi_3, \xi_4) = (\eta_1 \eta_2, \eta_2^{-1}, \eta_3, \eta_4). \quad (9)$$

Therefore, λ defines an isomorphism between X_{Ξ^*} and $\text{Hilb}^G(\mathbb{C}^4)$.

By (2), the canonical bundle of X_{Ξ^*} is given by $\omega_{X_{\Xi^*}} = \mathcal{O}_{X_{\Xi^*}}(E)$ with $E = D_c$. As the star of c in Ξ^* is given by the octahedron in [Fig. 1], we have $E \simeq (\mathbb{P}^1)^3$. One can apply the toric technique to determine the (-1) -hyperplane structure of fiber $\mathbb{P}^1$ of the (k, l) -th factor projection, $p_j : E \rightarrow (\mathbb{P}^1)^2$ for the normal bundle of E . For example, in the case of the projection of E onto $(\mathbb{P}^1)^2$ corresponding to the 2-convex set $\langle v^{12}, v^{13}, v^{34}, v^{24} \rangle$, the relation (9) between the local coordinates of X_{C_1} and $X_{C'_4}$ implies that the restriction of the normal bundle of E on each fiber $\mathbb{P}^1$ over $(3, 4)$ -plane is the (-1) -hyperplane bundle. This proves Theorem 4.1.

QED

By the property of the normal bundle of the canonical divisor E on $\mathbb{P}^1$ -fibers in Theorem 4.1, one can blow down E to obtain three different crepant resolutions of S_G . In fact, all these crepant resolutions are toric varieties described as follows: Let Ξ_j be the refinement of Ξ by adding the segment connecting v^{j4} and v^{kl} to divide the central polygon $\diamond$ into 4 simplices, where $\{j, k, l\} = \{1, 2, 3\}$. Then, we have the relation of refinements: $\Xi \prec \Xi_j \prec \Xi^*$ for $j = 1, 2, 3$, and each X_{Ξ_j} is a crepant resolution of X_{Ξ} . For the corresponding decompositions of the central core $\diamond$, the refinement relations are given by $\diamond \prec \diamond_j \prec \diamond^*$, ($1 \leq j \leq 3$), with the pictorial realization [Fig. 3]. By the relation between $X_{\diamond^*}$ and $X_{\diamond}$, one can conclude that $x_{\diamond}$ is the isolated singularity of $X_{\diamond}$, with the equation (5). The three 4-folds $X_{\diamond_j}$ are all “small”¹ resolutions of the singular variety $X_{\diamond}$. The relationship between these three crepant resolutions of the isolated singularity is defined to be the flop of 4-folds. Hence, we have shown the following result:


THEOREM 4.2 *There are three crepant resolutions of $S_{A_1(4)}$ obtained by blowing down the canonical divisor E of $\text{Hilb}^{A_1(4)}(\mathbb{C}^4)$ in Theorem 4.1. Any two such resolutions differ by a flop of 4-folds.*

Remark: The S_4 -action on coordinates leaves $\diamond$ and $\diamond^*$ invariant, but permutes three $\diamond_i$'s. A similar phenomenon will also appear in the case $G = A_1(5)$ in the next section.

5 $A_1(5)$ -Hilbert Scheme and Crepant Resolutions of $\mathbb{C}^5/A_1(5)$

In this section, we shall discuss the case of $G = A_1(5)$. Throughout this section, the indices i, j, k, l, m always mean a permutation of $(1, 2, 3, 4, 5)$. In addition to the elements e^i, v^{ij} of N in Δ , we consider

¹Here a “small” resolution means a resolution with the exceptional locus of codimension ≥ 2 .



[Fig. 3] Toric representation of flops of 4-folds over the isolated singularity and dominated by the smooth 4-fold.

the following rational points:

$$u^i = \frac{1}{4} \sum_{\alpha \neq i} e^\alpha, \quad w^i = \frac{1}{6} \left(e^i + \sum_{\alpha=1}^5 e^\alpha \right).$$

Then, one has a refinement Ξ^* of Ξ with elements in $\Xi^*(4)$ given by:

$$\Delta_i = \langle e^i, v^{ij}, v^{ik}, v^{il}, v^{im} \rangle, \quad \mathbf{I}_i = \langle w^i, v^{ij}, v^{ik}, v^{il}, v^{im} \rangle, \quad \mathbf{II}_{ij} = \langle w^i, u^j, v^{ik}, v^{il}, v^{im} \rangle,$$

$$\mathbf{III}_{jk} = \mathbf{III}_{kj} = \langle u^j, u^k, v^{im}, v^{lm}, v^{il} \rangle, \quad \mathbf{IV}_{i,jk} = \mathbf{IV}_{i,kj} = \langle w^i, u^j, u^k, v^{il}, v^{im} \rangle,$$

$$\mathbf{V}_{im} = \mathbf{V}_{mi} = \langle w^i, w^m, u^j, u^k, u^l, v^{im} \rangle, \quad \mathbf{VI} := \langle u^\alpha, w^\alpha \rangle_{\alpha=1}^5.$$

There are 5, 5, 20, 10, 30, 10, 1 elements of Δ_i , $\mathbf{I}_i$, $\mathbf{II}_{ij}$, $\mathbf{III}_{jk}$, $\mathbf{IV}_{i,jk}$, $\mathbf{V}_{im}$, $\mathbf{VI}$ respectively. The facet relations for these 4-polytopes can be depicted graphically in the following diagram:

$$\begin{array}{c} \Delta_i - \mathbf{I}_i - \mathbf{II}_{ij} \quad - \mathbf{IV}_{i,jk} - \mathbf{V}_{im} - \mathbf{VI} \\ | \\ \mathbf{III}_{jk} \end{array}$$

where a line describes the intersection of two polytopes with a common facet. Except the heptahedrons $\mathbf{V}_{im}$ and the decahedron $\mathbf{VI}$, the rest of the 4-polytopes are all simplicial.

THEOREM 5.1 *For $G = A_1(5)$, $\text{Hilb}^G(\mathbb{C}^5)$ is the toric variety X_{Ξ^*} with the canonical sheaf $\omega_{X_{\Xi^*}} = \mathcal{O}_{X_{\Xi^*}} \left(\sum_{i=1}^5 (2D_{w^i} + D_{u^i}) \right)$, where D_{u^i} and D_{w^i} are the divisors corresponding to u^i , w^i respectively.*

Proof: With (3) for $n = 5$, we have $U_i = \frac{Z_i}{Z_j Z_k Z_l Z_m}$, $V_{ij} = \frac{Z_i Z_j}{Z_k Z_l Z_m}$. The generators of $N \cap \sigma_{\mathcal{R}}$ and $M \cap \sigma_{\mathcal{R}}$ for $\mathcal{R} \in \Xi^*(4)$ are given by the following table:

$\mathcal{R}$	$N \cap \sigma_{\mathcal{R}}$	$M \cap \sigma_{\mathcal{R}}$
Δ_i	$e^i, v^{ij}, v^{ik}, v^{il}, v^{im}$	U_i, T_j, T_k, T_l, T_m
$\mathbf{I}_i$	$3w^i, v^{ij}, v^{ik}, v^{il}, v^{im}$	$U_i^{-1}, V_{ij}, V_{ik}, V_{il}, V_{im}$
$\mathbf{II}_{ij}$	$3w^i, 2u^j, v^{ik}, v^{il}, v^{im}$	$V_{ij}^{-1}, T_j, V_{ik}, V_{il}, V_{im}$
$\mathbf{III}_{jk}$	$2u^j, 2u^k, v^{im}, v^{lm}, v^{il}$	$T_j, T_k, V_{im}, V_{lm}, V_{il}$
$\mathbf{IV}_{i,jk}$	$3w^i, 2u^j, 2u^k, v^{il}, v^{im}$	$V_{lm}^{-1}, V_{ij}^{-1}, V_{ik}^{-1}, V_{il}, V_{im}$
$\mathbf{V}_{im}$	$3w^i, 3w^m, 2u^j, 2u^k, 2u^l, v^{im}$	$V_{ij}^{-1}, V_{jm}^{-1}, V_{ik}^{-1}, V_{km}^{-1}, V_{il}^{-1}, V_{lm}^{-1}, V_{im}$
$\mathbf{VI}$	$3w^i, 2u^i, (1 \leq i \leq 5)$	$V_{ij}^{-1}, (1 \leq i < j \leq 5)$

Then, one can easily see that $X_{\mathcal{R}}$ has the smooth toric structure $\mathbb{C}^5$ except $\mathcal{R} = \mathbf{V}_{im}$ and $\mathbf{VI}$, which contribute the singularities of X_{Ξ^*} . In fact, $X_{\mathbf{V}_{im}}$ is the affine variety in $\mathbb{C}^7$ defined by the relations,

$$V_{ij}^{-1} V_{km}^{-1} = V_{ik}^{-1} V_{jm}^{-1}, \quad V_{ik}^{-1} V_{lm}^{-1} = V_{il}^{-1} V_{km}^{-1}, \quad V_{il}^{-1} V_{jm}^{-1} = V_{ij}^{-1} V_{lm}^{-1}; \quad (10)$$

and $X_{\mathbf{VI}}$ is the affine variety in $\mathbb{C}^{10}$ given by

$$V_{ij}^{-1} V_{il}^{-1} V_{km}^{-1} = V_{ik}^{-1} V_{im}^{-1} V_{jl}^{-1}, \quad V_{ij}^{-1} V_{kl}^{-1} = V_{ik}^{-1} V_{jl}^{-1} \quad (11)$$

where the indices run through all possible i, j, k, l, m . For $\gamma \in \mathbb{C}$, we define the following eigen-polynomials of G :

$$F_{ij}(\gamma) := Z_i Z_j - \gamma Z_k Z_l Z_m, \quad H_{ij}(\gamma) := Z_k Z_l Z_m - \gamma Z_i Z_j.$$

We shall denote the coordinates of an element $y \in X_{\mathcal{R}}$ by

$\mathcal{R}$	$y \in X_{\mathcal{R}}$
Δ_i	$(U_i, T_j, T_k, T_l, T_m) = (\gamma_i, t_j, t_k, t_l, t_m)$
$\mathbf{I}_i$	$(U_i^{-1}, V_{ij}, V_{ik}, V_{il}, V_{im}) = (\gamma_{jklm}, \gamma_{ij}, \gamma_{ik}, \gamma_{il}, \gamma_{im})$
$\mathbf{II}_{ij}$	$(V_{ij}^{-1}, T_j, V_{ik}, V_{il}, V_{im}) = (\gamma_{klm}, t_j, \gamma_{ik}, \gamma_{il}, \gamma_{im})$
$\mathbf{III}_{jk}$	$(V_{im}, V_{il}, V_{lm}, T_k, T_j) = (\gamma_{im}, \gamma_{il}, \gamma_{lm}, t_k, t_j)$
$\mathbf{IV}_{i,jk}$	$(V_{il}, V_{im}, V_{ij}^{-1}, V_{ik}^{-1}, V_{lm}^{-1}) = (\gamma_{il}, \gamma_{im}, \gamma_{klm}, \gamma_{jlm}, \gamma_{ijk})$
$\mathbf{V}_{im}$	$(V_{im}, V_{il}^{-1}, V_{lm}^{-1}, V_{ik}^{-1}, V_{ij}^{-1}, V_{jm}^{-1}, V_{km}^{-1})$ $= (\gamma_{im}, \gamma_{jkm}, \gamma_{ijk}, \gamma_{jlm}, \gamma_{klm}, \gamma_{ikl}, \gamma_{ijl})$
$\mathbf{VI}$	$(V_{ij}^{-1})_{i < j} = (\gamma_{klm})_{k < l < m}$

where the γ 's and t 's are complex numbers, and in the cases of $\mathbf{V}_{im}$ and $\mathbf{VI}$, the coordinates are governed by the relations (10), (11) respectively. Using the above coordinates of y , one can express the ideal $I(y)$ as follows:

$$\begin{aligned}
\Delta_i \quad & I(y) = \langle Z_j^2 - t_j, Z_k^2 - t_k, Z_l^2 - t_l, Z_m^2 - t_m, Z_i - \gamma_i Z_j Z_k Z_l Z_m \rangle, \\
\mathbf{I}_i \quad & \begin{cases} I(y) = \langle Z_\alpha^2 - t_\alpha \rangle_{\alpha=1}^5 + \langle F_{ij}(\gamma_{ij}), F_{ik}(\gamma_{ik}), \\ F_{il}(\gamma_{il}), F_{im}(\gamma_{im}), Z_j Z_k Z_l Z_m - \gamma_{jklm} Z_i \rangle \\ \left(\begin{array}{l} t_j = \gamma_{ij} \gamma_{jklm}, \quad t_k = \gamma_{ik} \gamma_{jklm}, \quad t_l = \gamma_{il} \gamma_{jklm}, \\ t_m = \gamma_{im} \gamma_{jklm}, \quad t_i = \gamma_{ij} \gamma_{ik} \gamma_{il} \gamma_{im} \gamma_{jklm}^2 \end{array} \right) \end{cases} \\
\mathbf{II}_{ij} \quad & \begin{cases} I(y) = \langle Z_\alpha^2 - t_\alpha \rangle_{\alpha=1}^5 + \langle F_{ik}(\gamma_{ik}), F_{il}(\gamma_{il}), F_{im}(\gamma_{im}), H_{ij}(\gamma_{klm}) \rangle \\ \left(\begin{array}{l} t_k = \gamma_{ik} \gamma_{klm} t_j, \quad t_l = \gamma_{il} \gamma_{klm} t_j, \quad t_m = \gamma_{im} \gamma_{klm} t_j, \\ t_i = \gamma_{ik} \gamma_{il} \gamma_{im} t_j^2 \gamma_{klm} \end{array} \right) \end{cases} \\
\mathbf{III}_{jk} \quad & \begin{cases} I(y) = \langle Z_\alpha^2 - t_\alpha \rangle_{\alpha=1}^5 + \langle F_{im}(\gamma_{im}), F_{il}(\gamma_{il}), F_{lm}(\gamma_{lm}) \rangle, \\ (t_i = \gamma_{im} \gamma_{il} t_k t_j, \quad t_l = \gamma_{il} \gamma_{lm} t_k t_j, \quad t_m = \gamma_{im} \gamma_{lm} t_k t_j) \end{cases} \\
\mathbf{IV}_{i,jk} \quad & \begin{cases} I(y) = \langle Z_\alpha^2 - t_\alpha \rangle_{\alpha=1}^5 + \langle F_{im}(\gamma_{im}), F_{il}(\gamma_{il}), \\ H_{ij}(\gamma_{klm}), H_{ik}(\gamma_{jlm}), H_{lm}(\gamma_{ijk}) \rangle \\ \left(\begin{array}{l} t_m = \gamma_{im} \gamma_{jlm} \gamma_{klm} \gamma_{ijk}, \quad t_l = \gamma_{il} \gamma_{jlm} \gamma_{klm} \gamma_{ijk}, \\ t_i = \gamma_{im} \gamma_{il} \gamma_{jlm} \gamma_{klm} \gamma_{ijk}^2, \quad t_j = \gamma_{jlm} \gamma_{ijk}, \quad t_k = \gamma_{lmk} \gamma_{ijk} \end{array} \right) \end{cases} \\
\mathbf{V}_{im} \quad & \begin{cases} I(y) = \langle Z_\alpha^2 - t_\alpha \rangle_{\alpha=1}^5 + \langle F_{im}(\gamma_{im}), H_{il}(\gamma_{jkm}), H_{ik}(\gamma_{jlm}), \\ H_{ij}(\gamma_{klm}), H_{jm}(\gamma_{ikl}), H_{lm}(\gamma_{ijk}), H_{km}(\gamma_{ijl}) \rangle \\ \left(\begin{array}{l} t_i = \gamma_{im} \gamma_{ikl} \gamma_{jlm} \gamma_{ijk} = \gamma_{im} \gamma_{ikl} \gamma_{jkm} \gamma_{ijl}, \\ t_m = \gamma_{im} \gamma_{klm} \gamma_{jlm} \gamma_{ijk} = \gamma_{im} \gamma_{klm} \gamma_{jkm} \gamma_{ijl}, \\ t_l = \gamma_{ikl} \gamma_{jlm} = \gamma_{ijl} \gamma_{klm}, \quad t_j = \gamma_{ijk} \gamma_{jlm} = \gamma_{ijl} \gamma_{jkm}, \\ t_k = \gamma_{ijk} \gamma_{klm} = \gamma_{ikl} \gamma_{jkm} \end{array} \right) \end{cases} \\
\mathbf{VI} \quad & \begin{cases} I(y) = \langle Z_\alpha^2 - t_\alpha \rangle_{\alpha=1}^5 + \langle H_{lm}(\gamma_{ijk}) \rangle \text{ for all possible } i, j, k, l, m, \\ \left(\begin{array}{l} t_i = \gamma_{ijk} \gamma_{ilm} = \gamma_{ijm} \gamma_{ilk}, \gamma_{klm} \gamma_{jkm} \gamma_{ijl} = \gamma_{jlm} \gamma_{jkl} \gamma_{ikm} \\ \text{for all possible } i, j, k, l, m \end{array} \right) \end{cases}
\end{aligned}$$

One can show that the generators in the above expressions of $I(y)$ form the corresponding reduced Gröbner basis with respect to a weight order $\prec$ with weight $\mathbf{w} \in \text{Interior}(\sigma_{\mathcal{R}})$. Furthermore, the ideal $\text{lt}_{\prec}(I(y))$ is equal to $I(x_{\mathcal{R}})$. Therefore, $I(x_{\mathcal{R}})^\perp$ gives rise to a basis of $\mathbb{C}[Z]/I(y)$, which is G -regular by the following explicit description of basis elements.

$\mathcal{R}$	$I(x_{\mathcal{R}})^\perp$
Δ_i	$1, Z_j, Z_k, Z_l, Z_m, Z_j Z_k, Z_j Z_l, Z_k Z_l, Z_k Z_m, Z_l Z_m, Z_j Z_m,$ $Z_j Z_k Z_l, Z_j Z_k Z_m, Z_j Z_l Z_m, Z_k Z_l Z_m, Z_j Z_k Z_l Z_m$
$\mathbf{I}_i$	$1, Z_i, Z_j, Z_k, Z_l, Z_m, Z_j Z_k, Z_j Z_l, Z_k Z_l, Z_k Z_m, Z_l Z_m, Z_j Z_m,$ $Z_j Z_k Z_l, Z_j Z_k Z_m, Z_j Z_l Z_m, Z_k Z_l Z_m$
$\mathbf{II}_{ij}$	$1, Z_i, Z_j, Z_k, Z_l, Z_m, Z_i Z_j, Z_j Z_k, Z_j Z_l, Z_k Z_l, Z_k Z_m, Z_l Z_m,$ $Z_j Z_m, Z_j Z_k Z_l, Z_j Z_k Z_m, Z_j Z_l Z_m$
$\mathbf{III}_{jk}$	$1, Z_i, Z_j, Z_k, Z_l, Z_m, Z_i Z_j, Z_j Z_k, Z_j Z_l, Z_k Z_l, Z_k Z_m, Z_i Z_j Z_k,$ $Z_j Z_m, Z_j Z_k Z_l, Z_j Z_k Z_m, Z_i Z_k$
$\mathbf{IV}_{i,jk}$	$1, Z_i, Z_j, Z_k, Z_l, Z_m, Z_i Z_j, Z_j Z_k, Z_j Z_l, Z_k Z_l, Z_k Z_m, Z_l Z_m,$ $Z_j Z_m, Z_j Z_k Z_l, Z_j Z_k Z_m, Z_i Z_k$
$\mathbf{V}_{im}$	$1, Z_i, Z_j, Z_k, Z_l, Z_m, Z_i Z_j, Z_j Z_k, Z_j Z_l, Z_k Z_l, Z_k Z_m, Z_i Z_j Z_k,$ $Z_j Z_m, Z_j Z_k Z_l, Z_i Z_l, Z_i Z_k$
$\mathbf{VI}$	$1, Z_1, Z_2, Z_3, Z_4, Z_5, Z_1 Z_2, Z_1 Z_3, Z_1 Z_4, Z_1 Z_5, Z_2 Z_3, Z_2 Z_4,$ $Z_2 Z_5, Z_3 Z_4, Z_3 Z_5, Z_4 Z_5$

Thus, $\mathbb{C}[Z]/I(y)$ is a regular G -module for each $y \in X_{\Xi^*}$, and X_{Ξ^*} is birational over S_G having the vector bundle $\mathcal{F}_{\Xi^*}$. Hence, there is an epimorphism $\lambda : X_{\Xi^*} \rightarrow \text{Hilb}^G(\mathbb{C}^5)$ with $I(\lambda(y)) = I(y)$. With

the same argument as in the case $n = 4$, one can show that λ is injective on each $X_{\mathcal{R}}$ by the reason that the toric coordinates are encoded in the generators of $I(y)$. By the toric variety structure of X_{Ξ^*} , one obtains the injectivity of λ . Hence $X_{\Xi^*} \simeq \text{Hilb}^G(\mathbb{C}^5)$. By (2), one has the expression of the canonical sheaf for X_{Ξ^*} .

QED.


We are going to construct certain crepant resolutions of S_G dominated by the toric variety $X_{\Xi^*} \simeq \text{Hilb}^G(\mathbb{C}^5)$. Consider the permutation $\tau = (12345)$ on the coordinates of N_0 . Define the following simplices in Δ :

$$C := \langle v^{12}, v^{23}, v^{34}, v^{45}, v^{15} \rangle,$$

$$D_0 := \langle v^{12}, v^{23}, v^{34}, v^{13}, v^{15} \rangle,$$


$$E_0 := \langle v^{35}, v^{23}, v^{25}, v^{45}, v^{15} \rangle,$$

and $D_i := \tau^i(D_0)$, $E_i := \tau^i(E_0)$ ($i = 1, \dots, 4$). Then, one can show that C , D_i and E_i ($0 \leq i \leq 4$) form an integral simplicial decomposition of the central core $\diamond \in \Xi(4)$, hence one has the refinement $\diamond'$ of $\diamond$ with $\diamond'(4) = \{C\} \cup \{D_i, E_i\}_{i=0}^4$. Denote Ξ' the rational polytope decomposition of Δ with $\Xi'(4) = \diamond'(4) \cup \{\Delta_i\}_{i=1}^5$. Then Ξ' is a refinement of Ξ . By computing the primitive generators of each element in $\Xi'(4)$, one can easily check that $X_{\Xi'}$ is a crepant resolution of S_G . The facet relations of simplices in $\Xi'(4)$ are given in the following diagram:




LEMMA 5.1 *We have the relation $\diamond \prec \diamond' \prec \diamond^*$, consequently, $\Xi \prec \Xi' \prec \Xi^*$.*

Proof: It suffices to show the relation $\diamond' \prec \diamond^*$. By a detailed analysis of the 4-polytopes in $\Xi^*(4)$, one can conclude that the simplex C is the union of the following eleven polytopes in $\Xi'(4)$ with the facet relations:



Similarly, the relations of D_0, E_0 with 4-polytopes in $\diamond^*(4)$ are given in the following diagram:



By applying τ^i ($1 \leq i \leq 4$) on the above diagrams for D_0 and E_0 , one obtains the decompositions of D_i, E_i in terms of elements in $\diamond^*(4)$. The results then follow.

QED

As known in §2, Ξ and Ξ^* are S_5 -invariant polytope decompositions of Δ , but not for Ξ' . The isotropy subgroup of S_5 for Ξ' is $G' = \langle (25)(34), (12345) \rangle$, which is isomorphic to the dihedral group of order 10, with the index $[S_5 : G'] = 12$. By applying permutation elements in S_5 to the decomposition Ξ' , one obtains twelve integral simplicial refinements Ξ_i ($1 \leq i \leq 12$) of Ξ , with $\Xi \prec \Xi_i \prec \Xi^*$. Correspondingly, there are twelve decompositions $\diamond_i$ ($1 \leq i \leq 12$) of the central core $\diamond$ with the following refinement relations:

$$\diamond \prec \diamond_i \prec \diamond^*, \quad i = 1, \dots, 12.$$

The connection between these twelve smooth 5-folds corresponding to the toric data $\diamond_i$'s can be regarded as the “flop” of 5-folds, all of which are crepant resolutions of the singular variety defined by (6). Therefore, we have obtained the following result:

THEOREM 5.2 *There are twelve toric crepant resolutions, X_{Ξ_i} $1 \leq i \leq 12$, of $S_{A_1(5)}$ which are dominated by $\text{Hilb}^{A_1(5)}(\mathbb{C}^5)$. Any two such resolutions differ by a “flop” of 5-folds.*

References

- [1] E. Brieskorn, Rationale singularitäten komplexer Flächen, *Invent. Math.* 4 (1967) 336-358.
- [2] T. Bridgeland, A. King and M. Reid, The McKay correspondence as an equivalence of derived categories, *J. Amer. Math. Soc.* **14**(2001), no. 3, 535-554.
- [3] L. Chiang and S. S. Roan, Orbifolds and finite group representations, *Internat. J. of Math. and Math. Sci.* 26:11(2001)649-669. [math.AG/0007072](#); On Hypersurface quotient singularity of dimension 4. [math.AG/0011151](#).
- [4] D. Cox, J. Little and D. O'Shea, *Ideals, Varieties, and Algorithms*, Springer-Verlag, New York-Berlin-Heidelberg, 1992.
- [5] L. Dixon, J. Harvey, C. Vafa and E. Witten, Strings on orbifolds, *Nucl. Phys. B* 261 (1985) 678-686, Strings on orbifolds II, *Nucl. Phys. B* 274 (1986) 285-314.
- [6] F. Hirzebruch and T. Höfer, On the Euler number of an orbifold, *Math. Ann.* 286 (1990) 255-260.
- [7] Y. Ito, Crepant resolution of trihedral singularities. *Proc. Japan Acad. Ser. A, Math. Sci.* 70 (1994), no. 5, 131-136.
- [8] Y. Ito and H. Nakajima, McKay correspondence and Hilbert schemes in dimension three, *Topology* 39 (2000) 1155-1191, [math.AG/9803120](#).
- [9] Y. Ito and I. Nakamura, McKay correspondence and Hilbert schemes, *Proc. Japan Acad.* 72 (1996) 135-138.

- [10] G. Kempf, F. Knudson, D. Mumford and B. Saint-Donat, *Toroidal Embedding 1*, Lecture Notes in Math., Vol. 339, Springer-Verlag, New York, 1973.
T. Oda, *Lectures on Torus Embeddings and Applications*, Tata Inst. of Fund. Research 58, Springer-Verlag (1978).
V. I. Danilov, The geometry of toric varieties, Russ. Math. Surveys 33:2 (1978) 97-154.
- [11] F. Klein, *Gesammelte Mathematische Abhandlungen Bd. II*, Springer-Verlag, 1922 (reprint 1973).
- [12] D. G. Markushevich, M. A. Olshanetsky and A. M. Perelomov, Description of a class of superstring compactifications related to semi-simple Lie algebra, Comm. Math. Phys. 111 (1987) 247-274.
- [13] D. Markushevich, Resolution of $\mathbf{C}^3/H_{168}$, Math. Ann. 308 (1997) 279-289.
- [14] G. A. Miller, H. F. Blichfeldt and L. E. Dickson, *Theory and applications of finite groups*, John Wiley and Sons. Inc. (1916).
- [15] J. McKay, Graphs, singularities and finite groups, Proc. Symp. Pure Math. 37 (1980) 183-186.
- [16] I. Nakamura, Hilbert schemes of abelian group orbits, J. Alg. Geom. 10(4)(2001) 757-779.
- [17] S.S. Roan, Degeneration of K3 and abelian surfaces, Ph.D. thesis of Brandeis Univ. USA, 1974;
V. Kulikov, Degeneration of K3 and Enriques surface, Math. USSR Izvestija 11 (1977) 957-989;
U. Persson and H. Pinkham, Degeneration of surfaces with trivial canonical divisor, Ann. of Math. 113(1981) 45-66.
- [18] S. S. Roan and S. T. Yau, On Ricci flat 3-fold, Acta Math. Sinica, New Series 3 (3) (1987) 256-288.
- [19] S. S. Roan, On the generalization of Kummer surfaces, J. Diff. Geom. 30 (1989) 523-537.
- [20] S. S. Roan, On $c_1 = 0$ resolution of quotient singularity, Intern. J. Math. 5(4)(1994) 523-536.
- [21] S. S. Roan, Minimal resolutions of Gorenstein orbifolds in dimension three, Topology 35 (1996) 489-508.
- [22] I. Satake, On a generalization of the notion of manifold, Proceeding of National Academy Science, U.S.A. 42 (1956) 359-363.